



ISSN 2724-508X

<https://pagepress.org/medicine/acbr/index>

Publisher's disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

Annals of Clinical and Biomedical Research is, therefore, e-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one. The final version of the manuscript will then appear in a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

All legal disclaimers applicable to the journal apply to this production process as well.

Annals of Clinical and Biomedical Research 2023 [online ahead of print]

To cite this article:

I. B. Ndionuka, I. O. Onwuekwe, O. S. Ekenze, et al. Instantaneous stroke risk assessment of adults in Enugu, South East Nigeria: results from a random point survey. *Annals of Clinical and Biomedical Research*. 2023;4:206.
doi:10.4081/acbr.2023.206

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Instantaneous stroke risk assessment of adults in Enugu, South East Nigeria: results from a random point survey

Ijeoma. B. Ndionuka, Ikenna O. Onwuekwe, Oluchi S. Ekenze, Birinus Ezeala-Adikaibe, Chidinma Orah, Victor Madu, Chiamaka Okereke

Neurology Unit and Stroke Unit, Department of Medicine, University of Nigeria Teaching Hospital, Ituku-Ozalla, Enugu, Enugu State, Nigeria

Corresponding author: Ijeoma Ndionuka, Neurology and Stroke Unit, Department of Medicine, University of Nigeria Teaching Hospital, P.M.B. 01129 Enugu, Enugu State, Nigeria.

E-mail: ijeomandionuka@gmail.com

Key words: stroke risk, adults, Enugu, South East Nigeria.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Authors' contributions: all the authors made a substantive intellectual contribution. All the authors have read and approved the final version of the manuscript and agreed to be held accountable for all aspects of the work.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Abstract

Stroke remains a major cause of morbidity, death, and disability around the world, most especially in black Africans. Understanding the magnitude and appreciation of stroke risk in susceptible populations remains a key strategy in reducing stroke burden.

This study examined and compared 5-year stroke risk in an adult population in Enugu, Southeast Nigeria.

A point cross-sectional survey assessing stroke risk in adults was done in October 2019 in the city of Enugu as part of the World Stroke Day commemoration using the Stroke Riskometer app, and the data was analyzed. Ethical approval was obtained.

The total number of participants in this survey was 117 traders in a market in the Enugu metropolis, Enugu State, Nigeria. More males, 63%, than females, 37%, participated in this study. The total mean age was 47.50 years, with a total mean 5-year stroke risk of 3.93. Stroke risk was noted to be higher in females 4.56 than in males 3.56. Elevated blood pressure was present in 35% of the participants, and elevated blood glucose was seen in 5% of the total population.

The World Stroke Organization's Stroke Riskometer app is a valuable tool in assessing stroke risks especially in resource poor settings. This app also serves as a cost-effective method of conducting epidemiological research in stroke and other non-communicable diseases internationally.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Hypertension remains the most common risk factor for stroke globally. Higher stroke risk in females likely associated with higher mean age.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Introduction

The socioeconomic and health effects of stroke are huge and increasing worldwide, but especially so for developing countries and black populations. However, there is still low awareness about stroke and its risk factors in such Low- and Middle-Income Countries (LMICs). These factors, coupled with the under-utilization of strategies for the primary prevention of stroke and the lack of accurate data on the prevalence and effect of risk factors in different countries and populations, have been implicated in the ever-increasing worldwide burden of Non-Communicable Diseases (NCDs) including stroke.¹⁻⁴

The burden of stroke has also increased in young and middle-aged adults <65 years, which can be related to the increase in overweight/obesity caused by high consumption of fat diet. Knowledge about the contribution of modifiable risk factors to the increasing global and regional burdens of stroke is crucial for informing stroke prevention strategies.

The current prevalence of stroke in Nigeria is 1.14 per 1000, while the 30-day case fatality rate is as high as 40%.⁵ Management of the disease is largely conservative, while there is little or no funding for high-quality research.⁵

A crude stroke prevalence rate of 1.31/1000 population was observed in three semi-urban communities in the middle belt region of Nigeria. The prevalence of stroke was higher among the males than the females (1.54/1000 vs 1.08/1000) with a ratio of 1.4:1. Sixteen subjects (94.1%)

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

had one or more risk factors for stroke. Uncontrolled systemic hypertension (82.4%) was the most common risk factor for stroke, followed by Transient Ischemic Attack (TIA) (41.2%).⁶

There is a paucity of literature and data on stroke risk as assessed in the South East zone of Nigeria, of which Enugu serves as the capital. This study served to reflect the scope and magnitude of the stroke risk debacle in the south eastern region of Nigeria and to serve as a basis for greater concerted efforts to be made.

This paper seeks to assess stroke risks in selected populations of Nigerian adults utilizing the World Stroke Organization (WSO) Stroke Riskometer app which has been demonstrated to be a simple and yet efficient instrument for predicting stroke risk. The results reflect still the poor awareness among the general population and a need for the implementation of new, effective, widely available and cost-effective prevention and treatment strategies to reduce the incidence and severity of stroke in Nigeria.

Materials and Methods

Study site

Enugu state, from the 2006 National census in Nigeria, had a population of 3,267,837 people and was projected to have an estimated population of over 4.4M as of 2016.⁷ The state lies

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

geographically between longitudes 7^o 26^o E to 30^o E and latitudes 6^o 23^o N to 28^o N. The capital is Enugu City, with a population of 722,664 according to the 2006 Nigerian census.

Study design

This was a descriptive cross-sectional community survey of stroke risk in adults seen in a big market in Enugu metropolis, South East Nigeria.

Study population, resources and size

One hundred and seventeen adults aged >18 years were recruited during a public health enlightenment campaign as part of the commemoration of World Stroke Day by the Neurology and Stroke Units of the University of Nigeria Teaching Hospital Enugu, on October 29, 2019, in a market in Enugu metropolis. Consenting adults had their sociodemographic data obtained and their 5-year cumulative stroke risk assessed using the WSO Stroke Riskometer app. The validated Stroke Riskometer app provides the stroke probability score comparable to 2 well-known stroke prediction tools - the Framingham Stroke Risk Score and QSTROKE stroke risk prevention algorithms - commonly used in the medical field.^{8,9} The Stroke Riskometer app involves a high degree of interactivity and personalization and has received endorsement from the World Federation of Neurology, the World Stroke Organization, and the World Heart Federation.¹⁰ This

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

app has been demonstrated to be a simple, cheap, easily accessible, and effective instrument for determining the risk of stroke in an individual for the next 5 years.¹⁰

Results

The total number of participants in this study was 117 traders in a market in Enugu metropolis, Enugu State, Nigeria (*Table 1*).

More males, 63%, than females, 37%, participated in this study. The total mean age was 47.50 years, with a total mean stroke risk of 3.93. Stroke risk was noted to be higher in females, 4.56 (*Table 2*). The number of participants with elevated blood pressure was 36% of males and 33% of females. The number of participants with elevated blood glucose was 7% for males and 2% for females (*Table 3*).

Discussion

The recent INTERSTROKE case-control study, conducted in 22 countries worldwide, provided evidence that, collectively, 10 risk factors accounted for 88.1% (99% CI, 82.3%-92.2%) of the population-attributable risk for all stroke.¹¹ Because many (but not all) of these risk factors are modifiable, the INTERSTROKE data suggest that interventions that reduce blood pressure,

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

promote physical activity, smoking cessation, and a healthy diet, could substantially reduce the burden of stroke.

The Global Burden of Diseases 2010 studies found that the 3 leading risk factors for disease burden were high blood pressure, tobacco smoking (including second-hand smoke), and alcohol use.¹² Moreover, these 3 risk factors were ranked in the top 3 for several low- to middle-income regions, including Central and Eastern Europe, Southern Latin America, and Central Asia. In addition, lifestyle factors, such as high sodium, low fruit diets, and physical inactivity, contributed to 10% of disability-adjusted life years in 2010. These risk factors, along with the nonmodifiable factors of age, sex, and ethnicity, contribute significantly to stroke incidence.¹¹

Feigin *et al.*¹³ observed globally, 90.5% (95% UI 88.5-92.2) of the stroke burden (as measured in DALYs) was attributable to the modifiable risk factors analyzed, including 74.2% (95% UI 70.7-76.7) due to behavioral factors (smoking, poor diet, and low physical activity).

The Stroke Riskometer App is an important tool in spreading awareness about stroke, its risk factors, and how to tackle them. The Stroke Riskometer revealed a higher stroke risk in females among traders in the Enugu metropolis. This trend was also seen in global reports or studies on stroke risk assessments. The American Stroke Association reported that women have a higher stroke risk than men.¹⁴ Several factors like pregnancy, preeclampsia, birth control pills, and hormonal replacements were implicated in the higher stroke risk in females.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

In this risk assessment conducted among traders in the Enugu metropolis, elevated blood pressure contributed more to stroke risk than elevated blood glucose. In sub-Saharan Africa, Nigeria has the highest number of people with diabetes, with an estimated 3.9 million people (or an extrapolated prevalence of 4.99%) of the adult population aged 20-79 years old.¹⁵ This pattern was also observed in this study, with a total frequency of elevated blood glucose of 5%. The frequency of elevated blood pressure in this survey was 36% in males and 33% in females, which reflects the higher proportion of hypertension in males as seen in several studies done in Nigeria. Adeloje *et al.* noted an estimate of 29.5% in males and 25% in females in a meta-analysis of the prevalence of hypertension in Nigeria.¹⁶

Conclusions

The WSO Stroke Riskometer App is a valuable tool in assessing stroke risks, especially in resource-poor settings. This app also serves as a cost-effective method of conducting epidemiological research on stroke and other non-communicable diseases internationally.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

References

1. Strong K, Mathers C, Bonita R. Preventing stroke: saving lives around the world. *Lancet Neurol.* 2007;6:182-7.
2. Kontis V, Mathers CD, Rehm J, et al. Contribution of six risk factors to achieving the 25×25 non-communicable disease mortality reduction target: a modelling study. *Lancet.* 2014;384:427-37.
3. Sacco RL, Smith SC, Holmes D, et al. Accelerating progress on non-communicable diseases. *Lancet.* 2013;382:e4-5.
4. Feigin VL, Norrving B. A new paradigm for primary prevention strategy in people with elevated risk of stroke. *Int J Stroke.* 2014;9:624-6.
5. Wahab KW. The burden of stroke in Nigeria. *Int J Stroke.* 2008;3:290-2.
- 6.. Sanya EO, Desalu OO, Adepoju F, et al. Prevalence of stroke in three semi-urban communities in middle-belt region of Nigeria: a door to door survey. *Pan African Medical Journal.* 2015;20:33.
7. Nigeria Population Census. 2006. Available from: <https://nigeria.opendataforafrica.org/xspplpb/nigeria-census>
8. Parmar P, Krishnamurthi R, Ikram MA, et al. The Stroke Riskometer TM app: validation of a data collection tool and stroke risk predictor. *Int J Stroke.* 2015;10:231-44.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

9. Feigin VL, Krishnamurthi R, Bhattacharjee R, et al. New strategy to reduce the global burden of stroke. *Stroke*. 2015;46:1740-7.
10. World Stroke Organization. Stroke Riskometer. Available from: <http://www.strokeriskometer.com/>
11. O'Donnell MJ, Xavier D, Liu L, et al. INTERSTROKE investigators. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet*. 2010;376:112-23.
12. Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224-60.
13. Feigin VL, Roth GA, Naghavi M, et al. Global Burden of Diseases, Injuries and Risk Factors Study 2013 and Stroke Experts Writing Group. Global burden of stroke and risk factors in 188 countries, during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet Neurol*. 2016;15:913-24.
14. American Stroke Association. Women have a higher risk of stroke. 2018. Available from: <https://www.stroke.org/en/about-stroke/stroke-risk-factors/women-have-a-higher-risk-of-stroke>

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

15. World Health Organization. Follow-up to the Political Declaration of the High-level Meeting of the General Assembly on the Prevention and Control of Non-communicable Diseases. 2013.

Available from: <https://apps.who.int/iris/handle/10665/150161>

16. Adeloje D, Basquill C, Aderemi AV, et al. An estimate of the prevalence of hypertension in Nigeria: a systematic review and meta-analysis. *J Hypertens*. 2015;33:230-42.

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Table 1. Study population.

Sex	Number of participants	Mean age (years)	Modal age
Male	74	46.73	45
Female	43	48.81	52
Total	117	47.50	53

Table 2. Percentage of stroke risks.

Sex	Number of participants	Mean 5-year risk	Modal 5-year risk
Male	74	3.56	2.5
Female	43	4.56	2.6
Total	117	3.93	2.6

Table 3. Population with Elevated Blood Pressure (EBP)/Elevated Blood Glucose (EBG).

Sex	EBP	EBG	Total
Male	27(36%)	5(7%)	32

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Female	14(33%)	1(2%)	15
Total	41(35%)	6(5%)	47

The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.